


**Assessment Schedule – 2015****Chemistry: Demonstrate understanding of bonding, structure, properties and energy changes (91164)****Evidence Statement**

Q	Evidence	Achievement	Achievement with Merit	Achievement with Excellence
ONE (a)	$\begin{array}{c} \cdot\cdot \\ \text{O}=\text{O} \\ \cdot\cdot \end{array}$ $\begin{array}{c} \cdot\cdot \\ \text{Cl}-\text{O}-\text{Cl} \\ \cdot\cdot \end{array}$ $\begin{array}{c} \cdot\cdot \\ \text{O} \\ \parallel \\ \text{H}-\text{C}-\text{H} \end{array}$	<ul style="list-style-type: none"> <li>Two Lewis structures OR electron dot diagrams correct.</li> </ul>		
(b)	<p>In each <math>\text{CCl}_4</math> molecule, there are four negative / electron : densities / clouds / regions around the central C atom. These repel each other / are positioned as far away from each other as possible in a tetrahedral (base) arrangement, resulting in a <math>109.5^\circ</math> bond angle. All of these regions of electrons / electron densities are bonding, without any non-bonding regions, so the shape of the molecule is tetrahedral.</p> <p>In each <math>\text{COCl}_2</math> molecule, there are three negative / electron : densities / clouds / regions around the central C atom. These repel / are positioned as far away from each other as possible in a triangular / trigonal planar (base) shape, resulting in a <math>120^\circ</math> bond angle. All of these regions of electrons / electron densities are bonding, without any non-bonding regions, so the shape of the molecule is trigonal planar.</p>	<ul style="list-style-type: none"> <li>One shape with matching bond angle correct. OR Correctly identifies the number of electron densities surrounding the central atom of one molecule. OR States that the shape of the molecule is determined by the repulsion between regions of electron density around the central atom.</li> </ul>	<ul style="list-style-type: none"> <li>Links the shape of <u>both</u> molecules to the electron arrangement around the central atom. OR Links the bond angles in BOTH molecules to the electron arrangement around the central atom. OR Complete answer for <math>\text{CCl}_4</math> or <math>\text{COCl}_2</math>.</li> </ul>	<ul style="list-style-type: none"> <li>Evaluates the arrangement of electron densities around the central atom of BOTH molecules in order to correctly explain the shapes and bond angles.</li> </ul>

(c)	<p>Both molecules are non-polar.</p> <p>The Be-Cl bond is polar because Cl is more electronegative than Be / the atoms have different electronegativities.</p> <p>Since both the bonds are the same and arranged symmetrically around the central atom, in a linear arrangement, the bond dipoles cancel out, resulting in a non-polar molecule.</p> <p>The B-F bond is polar because F is more electronegative than B / the atoms have different electronegativities. Since all three bonds are the same and arranged symmetrically around the central atom, in a trigonal planar arrangement, the bond dipoles cancel out, resulting in another non-polar molecule.</p>				<ul style="list-style-type: none"><li>Identifies electronegativity difference between atoms.</li><li>OR</li><li>Identifies the polarity of either the Be-Cl or B-F bond correctly.</li><li>OR</li><li>States that polarity of the molecule depends on the symmetry of the molecule.</li></ul>	<ul style="list-style-type: none"><li>Non polar circled PLUS</li><li>Links the polarity of either Be-Cl or B-F bonds to the differences in electronegativity of the atoms involved.</li><li>OR</li><li>Links the even spread of polar bonds / bond dipoles around the central atom to their cancelling out and therefore to the overall non-polarity of the molecule.</li></ul>	<ul style="list-style-type: none"><li>Justifies choice of polarity in terms of polarity of bonds, due to differences in electro negativities of the atoms, and the cancelling out of bond dipoles / polar bonds due to the symmetry and shape of each molecule.</li></ul>												
(d)	<p><math>\text{C}_2\text{H}_4(\text{g}) + \text{Br}_2(\text{g}) \rightarrow \text{C}_2\text{H}_4\text{Br}_2(\text{g})</math></p> <table><tr><td><u>Bonds broken</u></td><td></td><td><u>Bonds formed</u></td></tr><tr><td>C=C      614</td><td></td><td>C–C      346</td></tr><tr><td>Br–Br      <u>193</u></td><td></td><td>C–Br      <u>2 × 285</u></td></tr><tr><td>807</td><td></td><td>916</td></tr></table> <p><math>\Delta_r H^\circ = \Sigma \text{Bond energies(bonds broken)} - \Sigma \text{Bond energies(bonds formed)}</math> <math>= 807 - 916 \quad (\text{or } 2463 - 2572)</math> <math>= -109 \text{ kJ mol}^{-1}</math></p> <p><i>(Alternative calculation that includes the breaking and reforming of four C-H bonds will also be accepted to Excellence level.)</i></p>							<u>Bonds broken</u>		<u>Bonds formed</u>	C=C      614		C–C      346	Br–Br <u>193</u>		C–Br <u>2 × 285</u>	807		916
<u>Bonds broken</u>		<u>Bonds formed</u>																	
C=C      614		C–C      346																	
Br–Br <u>193</u>		C–Br <u>2 × 285</u>																	
807		916																	
NØ	N1	N2	A3	A4	M5	M6	E7	E8											
No response or no relevant evidence.	1a	2a	3a	4a	2m	3m	2e	3e											

Q	Evidence	Achievement	Achievement with Merit	Achievement with Excellence
TWO(a)	Exothermic because the temperature of the solution increases / heat is released / particles slow down / bonds are formed	<ul style="list-style-type: none"> <li>One of (a) or (b)(i) is correctly identified with reason.</li> </ul>		
(b)(i)	Endothermic because the $\Delta_r H^\circ$ value is positive / it uses the sun's energy			
(ii)	$n(\text{CO}_2) = \frac{m}{M}$ $= \frac{19.8}{44.0}$ $= 0.450 \text{ mol}$ <p>Since 6 moles of <math>\text{CO}_2</math> reacting requires 2803 kJ of energy  then 1 mole of <math>\text{CO}_2</math> reacting requires <math>\frac{2803}{6} = 467.2</math> kJ of energy  and 0.450 moles of <math>\text{CO}_2</math> requires <math>467.2 \times 0.450 = 210</math> kJ of energy absorbed.</p>			
(c)(i)	$n(\text{C}_4\text{H}_{10}) = \frac{3.65}{58.0}$ $= 0.0629 \text{ mol}$ <p>If 0.0629 moles of <math>\text{C}_4\text{H}_{10}</math> releases 106 kJ of energy  Then 1 mole of <math>\text{C}_4\text{H}_{10}</math> releases <math>\frac{106}{0.0629} = 1685</math> kJ of energy  And 2 moles of <math>\text{C}_4\text{H}_{10}</math> releases <math>1685 \times 2 = 3370</math> kJ of energy (3368)  (<math>\Delta_r H = -3370 \text{ kJ mol}^{-1}</math>)</p>	<ul style="list-style-type: none"> <li>One step of calculation is correct.</li> </ul> OR Correct answer with no working.	<ul style="list-style-type: none"> <li>Two steps of the calculation for (b)(ii) are correct.</li> </ul>	<ul style="list-style-type: none"> <li>Calculation for (b)(ii) is correct with correct sign and units.</li> </ul>
		<ul style="list-style-type: none"> <li>One step of calculation is correct.</li> </ul> OR Correct answer with no working.	<ul style="list-style-type: none"> <li>Two steps of the calculation for (c)(i) are correct.</li> </ul>	<ul style="list-style-type: none"> <li>Calculation for (c)(i) is correct with correct sign and units.</li> </ul>

(ii)	<p>The results from this experiment are less than the accepted results, due to errors in the experimental design.</p> <p>The errors could include:</p> <ol style="list-style-type: none"> <li>1. Some energy is used to heat the metal can and the air surrounding the experiment / the experiment was not conducted in a closed system</li> <li>2. Incomplete combustion of butane.</li> <li>3. Some butane may have escaped before being ignited.</li> <li>4. The butane in the gas canister was impure.</li> <li>5. Some water evaporated</li> <li>6. Some energy was converted to light and sound</li> <li>7. Not carried out under standard conditions</li> </ol> <p>Therefore, not all of the energy released by the combustion of butane was transferred to heating the water.</p>	<ul style="list-style-type: none"> <li>Identifies an error. For example a user error, eg thermometer inaccurate. Measured water (volume) inaccurately. Scales inaccurate, for weighing butane canister.</li> </ul>	<ul style="list-style-type: none"> <li>Links a potential error in the experimental design, to why the actual results from the experiment were different to the accepted results.</li> </ul>	<ul style="list-style-type: none"> <li>Explains why the actual results from the experiment were less than the accepted results in relation to at least two experimental errors.</li> </ul>
(iii)		<ul style="list-style-type: none"> <li>Diagram correctly drawn, but not labelled.</li> </ul>	<ul style="list-style-type: none"> <li>Diagram correctly drawn and enthalpy change and reactants and products labelled correctly.</li> </ul> <p>AND</p> <p>Links overall enthalpy to bond-making releasing energy / being exothermic</p>	<ul style="list-style-type: none"> <li>Diagram correctly drawn and enthalpy change and reactants and products labelled correctly.</li> </ul> <p>AND</p> <p>Links overall enthalpy to BOTH bond-making and bond-breaking correctly.</p>

(iv)	When butane undergoes combustion, heat is released, so it is an exothermic reaction. Bond-making is an exothermic process / releases energy and bond-breaking is endothermic / requires energy. For the overall reaction in the combustion of butane to release energy, more energy is given out as bonds are made (when the products, CO <sub>2</sub> and H <sub>2</sub> O are formed) than the energy being used to break the bonds (in the reactants, C <sub>4</sub> H <sub>10</sub> and O <sub>2</sub> ).				<ul style="list-style-type: none"> <li>Identifies bond-making as exothermic / releases energy</li> </ul> OR <ul style="list-style-type: none"> <li>Bond-breaking as endothermic / absorbs or requires energy</li> </ul> OR <ul style="list-style-type: none"> <li>Bonds formed are stronger than bonds broken</li> </ul>			
NØ	N1	N2	A3	A4	M5	M6	E7	E8
No response or no relevant evidence	1a	2a	3a	4a	3m	4m	2e, including one explanation from part (c)	3e

Q	Evidence				Achievement	Achievement with Merit	Achievement with Excellence
THREE (a)	Substance	Type of Substance	Type of particle	Attractive forces between particles	• One row or one column correct.	• Three rows or two columns correct.	
	Cu(s)	metal / metallic	atom / cation and delocalised electrons / nuclei and electrons	metallic bond / electrostatic attraction between atom / cation / nuclei and electron			
	PCl <sub>3</sub> (s)	molecular	molecule	intermolecular (forces)			
	SiO <sub>2</sub> (s)	covalent network / giant covalent	atom	covalent bond			
	KCl(s)	ionic	ion	ionic bond			
(b)	Phosphorus trichloride, PCl <sub>3</sub> , is a molecular solid, made up of non-metal phosphorus and chlorine atoms covalently bonded together. The molecules are held together by weak intermolecular forces. Since these forces are weak, not much energy is required to overcome them, resulting in low melting / boiling points. (In the case of PCl <sub>3</sub> , its melting point is lower than, and its boiling point is higher than room temperature, so it is liquid.)  PCl <sub>3</sub> does not contain free moving ions nor any delocalised / free moving valence electrons, meaning PCl <sub>3</sub> does not contain any charged particles. Since free moving ions / electrons / charged particles are required to carry electrical current, PCl <sub>3</sub> is unable to conduct electricity.				• Reason given for one property of PCl <sub>3</sub> .	• Links either state or conductivity to structure and bonding for PCl <sub>3</sub> .	• Explanation links both state and conductivity to structure and bonding for PCl <sub>3</sub> .

(c)	<p>Cu is insoluble in water and malleable.</p> <p>Copper is a metal made up of an array of atoms / ions / nuclei held together by non-directional forces between the positive nuclei of the atoms and the delocalised / free moving valence electrons. There is no attraction between the copper atoms and the (polar) water molecules, therefore Cu is insoluble in water.</p> <p>Since the attractive forces are non-directional, when pressure is applied, the Cu atoms can move past each other to change shape without the bonds breaking, so Cu is malleable. (Note – labelled diagrams can provide replacement evidence).</p>				<ul style="list-style-type: none"><li>• Table completely correct.</li><li>• Reason given for malleability for any substance.</li><li>• Reason given for solubility for any substance.</li></ul>		<ul style="list-style-type: none"><li>• Links ONE property for ONE substance to its particles, structure, and bonding.</li><li>• Links ONE property for A SECOND substance to its particles, structure, and bonding.</li></ul>		<ul style="list-style-type: none"><li>• Justification links BOTH properties for ONE substance to its particles, structure, and bonding.</li><li>• Justification links BOTH properties for A SECOND substance to its particles, structure, and bonding.</li></ul>	
	<p>KCl is soluble in water and not malleable.</p> <p>KCl is made up of positive / <math>K^+</math> ions, and negative / <math>Cl^-</math> ions, ionically bonded in a 3D lattice. When added to water, polar water molecules form electrostatic attractions with the <math>K^+</math> and <math>Cl^-</math> ions. The partial negative charge, <math>\delta^-</math>, on oxygen atoms in water are attracted to the <math>K^+</math> ions and the partial positive, <math>\delta^+</math>, charges on the H's in water are attracted to the <math>Cl^-</math> ions, causing KCl to dissolve in water.</p> <p>KCl is not malleable because if pressure is applied to an ionic lattice, it forces ions with the same charge next to each other; they repel each other and break the structure. (Note – labelled diagrams can provide replacement evidence).</p>									
	<p><math>SiO_2</math> is insoluble in water and not malleable.</p> <p><math>SiO_2</math> is a covalent network made up of atoms covalently bonded together in a 3D lattice structure. (Covalent bonds are strong), Polar water molecules are not strong / insufficiently attracted to the Si and O atoms, therefore <math>SiO_2</math> is insoluble in water.</p> <p><math>SiO_2</math> is not malleable because if pressure is applied, the directional / strong covalent bonds have to be broken before the atoms can move.</p> <p>(Note - labelled diagrams can provide replacement evidence).</p>									
NØ	N1	N2	A3	A4	M5	M6	E7	E8		
No response or no relevant evidence	1a	2a	3a	4a	3m	4m	2e	3e		

### Cut Scores

Not Achieved	Achievement	Achievement with Merit	Achievement with Excellence
0 – 7	8 – 13	14 – 18	19 – 24