

No part of the candidate evidence in this exemplar material may be presented in an external assessment for the purpose of gaining credits towards an NCEA qualification.

2

91164



911640



NEW ZEALAND QUALIFICATIONS AUTHORITY
MANA TOHU MĀTAURANGA O AOTEAROA

QUALIFY FOR THE FUTURE WORLD
KIA NOHO TAKATŪ KI TŌ ĀMUA AO!

SUPERVISOR'S USE ONLY

Level 2 Chemistry, 2017

91164 Demonstrate understanding of bonding, structure, properties and energy changes

2.00 p.m. Thursday 16 November 2017
Credits: Five

Achievement	Achievement with Merit	Achievement with Excellence
Demonstrate understanding of bonding, structure, properties and energy changes.	Demonstrate in-depth understanding of bonding, structure, properties and energy changes.	Demonstrate comprehensive understanding of bonding, structure, properties and energy changes.

Check that the National Student Number (NSN) on your admission slip is the same as the number at the top of this page.

You should attempt ALL the questions in this booklet.

A periodic table is provided on the Resource Sheet L2-CHEMR.

If you need more room for any answer, use the extra space provided at the back of this booklet and clearly number the question.

Check that this booklet has pages 2–12 in the correct order and that none of these pages is blank.

YOU MUST HAND THIS BOOKLET TO THE SUPERVISOR AT THE END OF THE EXAMINATION.

Excellence

TOTAL

24

ASSESSOR'S USE ONLY

QUESTION ONE

- (a) When solid calcium chloride, $\text{CaCl}_2(s)$, reacts with water, the temperature increases.

Circle the term that best describes this reaction.

endothermic

exothermic

Give a reason for your choice.


An exothermic reaction releases energy eg. as heat, which increases the temperature of the surroundings.

- (b) When a person sweats, water is lost from the body by evaporation. This is an endothermic process. This evaporation speeds up when a person exercises.

- (i) Explain why the evaporation of water in sweat from the body is endothermic, and why exercise increases this evaporation.

Sweat evaporating is $\text{H}_2\text{O}(l) \rightarrow \text{H}_2\text{O}(g)$. This is an endothermic reaction as energy is required (absorbed from surroundings) to break the intermolecular attractions between H_2O molecules. When a

- (ii) Draw a labelled enthalpy diagram for the evaporation of water, $\text{H}_2\text{O}(l)$. surroundings



and evaporate to form $\text{H}_2\text{O}(g)$ at a greater rate.


- (iii) Sodium chloride, NaCl, is another compound that is excreted from the body in sweat.

Use your knowledge of structure and bonding to explain the dissolving process of sodium chloride, NaCl, in water.

Support your answer with a labelled diagram.


NaCl is an ionic solid with strong ionic bonds between Na^+ (cations) and Cl^- (anions). H_2O is a polar molecule with the O atom being slightly ~~pos~~ negative and the H atoms being slightly positive. When NaCl is dissolved in H_2O , the positive Na^+ are attracted to the δ^- O end of the H_2O molecule, and the negative Cl^- are attracted to the δ^+ (slightly positive) H end of the H_2O molecule as shown in the diagram. The attractions between Na^+ and H_2O , and Cl^- and H_2O are greater than the attractions between Na^+ and Cl^- , and H_2O and H_2O . Therefore, the ionic bond between Na^+ and Cl^- is ^{able to be} broken and Na^+ and Cl^- are surrounded by H_2O molecules. Thus, NaCl is soluble and dissolves in H_2O , water (polar solvent).

Space for diagram



- (c) Thermite reactions occur when a metal oxide reacts with a metal powder.

The equations for two thermite reactions are given below:



Use calculations to determine which metal oxide, iron(III) oxide, $\text{Fe}_2\text{O}_3(\text{s})$, or copper(II) oxide, $\text{CuO}(\text{s})$, will produce more heat energy when 50.0 g of each metal oxide is reacted with aluminium powder, $\text{Al}(\text{s})$.

$$M(\text{Fe}_2\text{O}_3) = 160 \text{ g mol}^{-1}$$

$$M(\text{CuO}) = 79.6 \text{ g mol}^{-1}$$

◦ Reaction 1:

$$n = \frac{m}{M} \quad n(\text{Fe}_2\text{O}_3) = \frac{50}{160} = 0.313 \text{ mol}$$



◦ Reaction 2:



$$n = \frac{m}{M} \quad n(\text{CuO}) = \frac{50}{79.6} = 0.628 \text{ mol}$$



Therefore, $\text{CuO}(\text{s})$ will produce ^(52 kJ) more heat energy when 50g of CuO is reacted with $\text{Al}(\text{s})$, as shown in reaction 2. //

QUESTION TWO

- (a) (i) Draw the Lewis structure (electron dot diagram) for the following molecules, and name their shapes.

Molecule	$\overset{16}{\text{O}} \overset{7}{\text{H}} \overset{17}{\text{Cl}} \quad 14$ HOCl	$\overset{6}{\text{C}} \overset{16}{\text{O}} \overset{17}{\text{Cl}} \quad 24$ COCl ₂	$\overset{7}{\text{N}} \overset{9}{\text{F}} \quad 21$ NF ₃ 26
Lewis structure			
Name of shape	bent	trigonal planar	trigonal pyramid.
Approximate bond angle around the central atom	109.5°	120°	109.5°

- (ii) Justify the shapes and bond angles of HOCl and COCl₂.

• HOCl has 4 regions of electron density around its central atom O. Since electrons repel and are maximum distance apart, HOCl has a bond angle of 109.5° (based on tetrahedral shape) around the central atom O. HOCl has 2 bonded and 2 non-bonded regions of electron density around central atom O, therefore has a bent shape as only the bonded regions are visible (bonded to H and Cl).

• COCl₂ has only 3 regions of electron density around its central atom C. Since electrons repel and are maximum distance apart, the bond angle around the central atom C is 120°. Since all 3 regions of electron density are bonded, the shape of COCl₂ is trigonal planar.

(b) Three-dimensional diagrams for two molecules are shown below.

Molecule		
Name	Dichloromethane	Tetrachloromethane
Polarity of molecule	polar	non-polar

- (i) In the boxes above, identify the polarity of each molecule, by writing either **polar** or **non-polar**.
- (ii) Justify your choices.



Dichloromethane has 2 bond dipoles from C to Cl and 2 ^(smaller) bond dipoles from H to C. There are bond dipoles between C and Cl as there is a difference in electronegativity (attracting pull of an atom on a bonded pair of electrons) \rightarrow Cl is more electronegative than C, so the electrons are closer to the Cl atom. There are also 2 smaller bond dipoles between C and H due to the difference in electronegativity

(C is more electronegative than H). Since the 4 bond dipoles around central atom C are unequally sized (asymmetrical), the bond dipoles do not cancel and thus dichloromethane is polar. ^(there is a molecular dipole)


Tetrachloromethane ~~also~~ has 4 equally sized bond dipoles ^(1 polar bonds) from C to Cl as Cl is more electronegative than C ^(electronegativity difference = bond dipole). Since these 4 bonds are arranged in a tetrahedral shape and are equally sized (symmetrical) the bond dipoles/polar bonds cancel out and tetrachloromethane is non-polar.

(c) Hydrazine, N_2H_4 , is used as rocket fuel.

Use calculations to determine which of **Reaction 1** or **Reaction 2** releases more energy.




The structure of each chemical species is shown in the box below.



Use the average bond enthalpies given in the table below.

Bond	Average Bond enthalpy /kJ mol ⁻¹	Bond	Average Bond enthalpy /kJ mol ⁻¹
H-H	436	N-N	158
H-F	567	F-F	159
N-H	391	O=O	498
O-H	463	N≡N	945

Show your working and include appropriate units in your answer.



Therefore reaction 2 releases more energy (per mol.)

QUESTION THREE

- (a) Complete the table below by stating the type of solid, the type of particle, and the type of bonding (attractive forces) between the particles in each solid.

Solid	Type of solid	Type of particle	Attractive forces between particles
Al(s) (aluminium)	metallic <u>lattice</u>	cations + <u>electrons</u>	metallic <u>bonds.</u>
MgCl ₂ (s) (magnesium chloride)	<u>ionic</u> lattice	cations + <u>anions</u>	<u>ionic</u> bonds
S ₈ (s) (sulfur)	<u>molecular</u> solid	<u>molecules</u>	(weak) intermolecular <u>forces</u>

- (b) Circle the substance which has the lowest melting point.

Al(s)

MgCl₂(s)S₈(s)

Justify your choice, referring to the attractive forces between the particles of ALL three substances.

Al(s) is a metallic lattice with strong non-directional metallic bonding between cations and a sea of delocalized electrons. MgCl₂ is an ionic lattice with strong ionic bonds between (Mg²⁺) cations and (Cl⁻) anions. S₈(s) is a molecule with weak intermolecular attractions to other S₈ molecules. Out of all 3 substances, S₈ has the weakest attractive forces between particles: weak intermolecular / vander waals forces. This means these attractions require the least energy to break compared to metallic bonds and ionic bonds. Therefore, S₈ has the lowest melting point as it requires the least (heat) energy / temperature (average measure of kinetic energy of particles) to weaken / breaks the attractive forces (weak intermolecular).

attractions) between particles (molecules).

ASSESSOR'S
USE ONLY

**Question Three
continues on the
following page.**

(c) Circle the substance which is malleable.

Al(s)

MgCl₂(s)

S₈(s)


Justify your choice by referring to the structure and bonding of your chosen substance.

You may include a diagram or diagrams in your answer.



Al(s) is a metallic lattice with strong non-directional metallic bonding between the lattice of cations (Al³⁺) and the sea of delocalized electrons. Due to this ^{strong} non-directional metallic bonding, Al(s) is malleable. When the metallic lattice is struck/hit, the layers of cations are able to slide past one another without disrupting the metallic bonds as the sea of delocalized electrons can freely move and maintain strong (non-directional) metallic bonds.

(However, MgCl₂ and S₈ are not malleable as their ionic bonds and weak intermolecular forces are easily broken when struck/hit as described in diagram below).

Space for diagram



NOT malleable



ES

**Extra paper if required.
Write the question number(s) if applicable.**

ASSESSOR'S
USE ONLY

QUESTION
NUMBER

The page contains a large grid of horizontal lines for writing answers. A vertical line is positioned on the left side, creating a column for question numbers. A diagonal red line is drawn across the grid from the bottom-left corner towards the top-right corner, indicating that the page is to be used as extra paper.

Extra paper if required.
Write the question number(s) if applicable.

ASSESSOR'S
USE ONLY

QUESTION
NUMBER

91164

Subject:	Chemistry	Standard:	91164	Total score:	24
Q	Grade score	Annotation			
1	E8	This candidate received a grade score of E8 as the response comprehensively relates increased heat from exercise to increased bond breaking during evaporation, evaluates the attractions of H ₂ O and NaCl with words and diagrams, as well as analysing and correctly contrasting the energy given out during thermochemical reactions.			
2	E8	This candidate received a grade score of E8 as the response elaborates fully on the factors that determine shape and polarity of molecules while also correctly evaluating the relative enthalpy changes of hydrazine reactions.			
3	E8	This candidate received a grade score of E8 as the response comprehensively relates bonding strength with melting points while also elaborating on the factors that enable the malleability of aluminium to occur.			