

## Assessment Schedule – 2017 V FINAL

## Chemistry: Demonstrate understanding of the properties of organic compounds (91391)



## Evidence Statement





Q	Evidence	Achievement	Merit	Excellence
ONE (a)  (b)	See <b>Appendix A</b> .  Reagent 1 = $\text{NaBH}_4 / \text{LiAlH}_4$ Reduction $\text{CH}_3 - \underset{\text{OH}}{\text{CH}} - \text{CH}_3$  Reagent 2 = conc. $\text{H}_2\text{SO}_4$ (heat) / $\text{Al}_2\text{O}_3$ Elimination $\text{H}_2\text{C} = \text{CH} - \text{CH}_3$  Addition $\text{CH}_3 - \underset{\text{Cl}}{\text{CH}} - \text{CH}_3$ $\text{CH}_3\text{CH}_2\text{CH}_2 - \text{Cl}$	<ul style="list-style-type: none"> <li>SIX correct.</li> <li>ONE correct reagent and one reaction type.</li> <li>TWO correct structures.</li> </ul>	<ul style="list-style-type: none"> <li>ALL correct.</li> <li>SEVEN correct.</li> </ul> OR All correct showing understanding of the chemistry but with repeated error.	<ul style="list-style-type: none"> <li>ALL NINE correct, including identification of both minor and major products.</li> </ul>
(c)(i)  (ii)	<p>There must be a carbon atom that has <b>four different species (groups)</b> attached to it. This creates two molecules that are <b>mirror images</b> of each other that are <b>non-superimposable</b>.</p> <p>The different isomers will <b>rotate (plane)-polarised light in opposite directions</b>. This will distinguish the isomers.</p>	Any of the following to a maximum of three: <ul style="list-style-type: none"> <li>One correct 3-D Drawing OR two 3-D drawings with the four correct groups</li> <li>States that four-different species are required</li> <li>Mirror images</li> <li>Non-superimposable</li> <li>Enantiomers will rotate plane-polarised light.</li> </ul>	<ul style="list-style-type: none"> <li>Correct 3-D drawings and partial explanation.</li> </ul> OR Full explanation with correct but careless drawings.	<ul style="list-style-type: none"> <li>Correct 3-D drawings with full explanation.</li> </ul>

<b>NØ</b>	<b>N1</b>	<b>N2</b>	<b>A3</b>	<b>A4</b>	<b>M5</b>	<b>M6</b>	<b>E7</b>	<b>E8</b>
No response; no relevant evidence.	1a	2a	3a	5a	2m	3m	1e + 2m	2e + 1m

Q	Evidence	Achievement	Merit	Excellence
TWO (a)	See <b>Appendix B</b> .	<ul style="list-style-type: none"> <li>• FOUR structures / reagents correct.</li> </ul>	<ul style="list-style-type: none"> <li>• SEVEN structures / reagents correct</li> <li>OR</li> <li>All correct showing understanding of the chemistry, but with repeated error.</li> </ul>	<ul style="list-style-type: none"> <li>• ALL correct.</li> </ul>
(b)(i)	<p>Aldehyde (propanal) is obtained by distillation of propan-1-ol with acidified (potassium) dichromate (<math>\text{Cr}_2\text{O}_7^{2-} / \text{H}^+</math>). The orange colour of the <math>\text{Cr}_2\text{O}_7^{2-} / \text{H}^+</math> changes to (blue) green (<math>\text{Cr}^{3+}</math> ions). The reaction is an oxidation reaction.</p> <p>Distillation is a way to separate the aldehyde (propanal) from the reactant alcohol (propan-1-ol) which has a higher boiling point. The aldehyde (propanal) can react further to form a carboxylic acid (propanoic acid). This reaction is prevented if the <b>aldehyde is removed as it is formed</b> – distillation achieves this by evaporating the aldehyde and then allowing it to condense for collection.</p>	<ul style="list-style-type: none"> <li>• Distillation identified.</li> <li>• Identifies oxidation reaction.</li> <li>• Correct colour change.</li> </ul>	<ul style="list-style-type: none"> <li>• Correct colour change, reaction type and recognises the need for distillation related to different boiling points / preventing further reaction.</li> <li>OR</li> <li>Correct answer, with one omission, e.g. oxidation or colour change.</li> </ul>	<ul style="list-style-type: none"> <li>• Full explanation of how only propanal is produced in the laboratory.</li> </ul>
(ii)	<p>Adding blue Benedict's solution to a <b>warmed / heated</b> sample of propanal will cause a (brick) red colour to form. This happens because the propanal has been oxidised to propanoic acid / carboxylic acid (red colour is copper(I) oxide).</p> $\text{CH}_3\text{CH}_2\underset{\text{O}}{\underset{\parallel}{\text{C}}}\text{H} + \text{Benedict's reagent} \xrightarrow{[\text{O}]} \text{CH}_3\text{CH}_2\underset{\text{O}}{\underset{\parallel}{\text{C}}}\text{OH}$ <p>No change will occur when blue Benedict's solution is added to propanone, as it cannot be further oxidised / won't react.</p>	<ul style="list-style-type: none"> <li>• Recognises that only propanal will react.</li> <li>• ONE of the following: EITHER correct colour change OR reaction type OR equation.</li> </ul>	<ul style="list-style-type: none"> <li>• Only propanal reacts, plus any TWO of: correct reaction type, correct colour change, correct equation.</li> </ul>	<ul style="list-style-type: none"> <li>• Explanation for both aldehyde and ketone, with correct equation.</li> </ul>

NØ	N1	N2	A3	A4	M5	M6	E7	E8
No response; no relevant evidence.	1a	2a	3a	5a	2m	3m	1e + 2m	2e + 1m

Q	Evidence	Achievement	Merit	Excellence
THREE (a)(i)	 <p>cysteine                      serine</p> <p>Second dipeptide the same structure above, with the CH<sub>2</sub>SH swapped with CH<sub>2</sub>OH.</p>	<ul style="list-style-type: none"> <li>• ONE correct dipeptide. OR Correct section of protein (a continuing structure) OR Both essentially correct, but includes careless errors.</li> </ul>	<ul style="list-style-type: none"> <li>• BOTH dipeptides correct.</li> </ul>	
(ii)	 <p>Amide linkage group circled on one of the dipeptides.</p>	<ul style="list-style-type: none"> <li>• Amide linkage group circled.</li> </ul>		

<p>(b)</p>	<p>Nomex® has an amide linkage – see (a)(ii) above.</p> <p>Monomers:</p> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p><chem>Nc1cccc(N)c1</chem></p> </div> <div style="text-align: center;">  <p><chem>OC(=O)c1ccc(C(=O)O)cc1</chem></p> </div> </div> <p>OR</p> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p><chem>Nc1ccc(N)cc1</chem></p> </div> <div style="text-align: center;">  <p><chem>ClC(=O)c1ccc(C(=O)Cl)cc1</chem></p> </div> </div> <p>This is a condensation polymer / polyamide, as monomers join / amide link forms and a molecule of water or HCl is released during the reaction.</p>	<ul style="list-style-type: none"> <li>• ONE correct monomer OR both correct but includes careless error.</li>   <li>• Amide functional group named and polymer, or reaction described as condensation or a polyamide</li>   <li>• States condensation reaction with some explanation.</li> </ul>	<ul style="list-style-type: none"> <li>• Both monomers correct.</li>   <li>• Condensation reaction fully explained.</li> </ul>	<ul style="list-style-type: none"> <li>• Full evaluation of Nomex®.</li> </ul>
------------	---	---	--	--

(c)  
(i)(ii)

Two forms of hydrolysis:

$\left[ \text{N}(\text{H})-\text{C}_6\text{H}_4-\text{N}(\text{H})-\text{C}(=\text{O})-\text{C}_6\text{H}_4-\text{C}(=\text{O}) \right]_n$

$\xrightarrow{\text{HCl/heat}}$   $\text{H}_3\text{N}^+-\text{C}_6\text{H}_4-\text{NH}_3^+$  +  $\text{HO}-\text{C}(=\text{O})-\text{C}_6\text{H}_4-\text{C}(=\text{O})-\text{OH}$

$\xrightarrow{\text{NaOH/heat}}$   $\text{H}_2\text{N}-\text{C}_6\text{H}_4-\text{NH}_2$  +  $\text{O}-\text{C}(=\text{O})-\text{C}_6\text{H}_4-\text{C}(=\text{O})-\text{O}$

- ONE structure correctly drawn in part (i).
- ONE structure correctly drawn in part (ii).

- ONE correct reaction. (part (i) or (ii) correct).
- OR
- All correct showing understanding of the chemistry, but with repeated error.

- BOTH reactions correct.

NØ	N1	N2	A3	A4	M5	M6	E7	E8
No response; no relevant evidence.	1a	2a	3a	5a	2m	3m	1e + 2m	2e + 1m

### Cut Scores

Not Achieved	Achievement	Achievement with Merit	Achievement with Excellence
0 – 7	8 – 14	15 – 19	20 – 24

**APPENDIX A.**

Question One (a)

*Boxes in grey are part of the question – answers are in white boxes.*

Functional group	Structural formula	IUPAC (systematic) name
Alkene	$\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2$	but-1-ene
Amine	$\begin{array}{c} \text{CH}_3\text{CHCH}_2-\text{N}-\text{H} \\   \quad   \\ \text{CH}_3 \quad \text{H} \end{array}$	2-methylpropan-1-amine
Acyl chloride	$\begin{array}{c} \text{CH}_3\text{CH}_2\text{CH}_2\text{C}-\text{Cl} \\    \\ \text{O} \end{array}$	butanoyl chloride 2-methylpropanoyl chloride
Ester	$\begin{array}{c} \text{H}-\text{C}-\text{O}-\text{CH}_2\text{CH}_2\text{CH}_3 \\    \\ \text{O} \end{array}$	propyl methanoate
Ketone	$\begin{array}{c} \text{CH}_3\text{CH}_2-\text{C}-\text{CH}_3 \\    \\ \text{O} \end{array}$	Butanone Butan-2-one
Aldehyde	$\begin{array}{c} \text{CH}_3\text{CH}_2\text{CH}_2-\text{C}-\text{H} \\    \\ \text{O} \end{array}$	Butanal 2-methylpropanal
Amide	$\begin{array}{c} \text{CH}_3\text{CH}_2\text{CH}_2-\text{C}-\text{NH}_2 \\    \\ \text{O} \end{array}$	butanamide

**APPENDIX B.**

## Question Two (a)

Compound	Structure
P	$\begin{array}{c} \text{CH}_3\text{CH}_2\text{CH}_2\text{CHCH}_3 \\   \\ \text{OH} \end{array}$
Q	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2 - \text{OH}$
R	$\text{CH}_3\text{CH} = \text{CHCH}_2\text{CH}_3$
S	$\text{CH}_2 = \text{CHCH}_2\text{CH}_2\text{CH}_3$
T	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2 - \text{Cl}$
U	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2 - \text{NH}_2$
V	$\begin{array}{c} \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{C} - \text{OH} \\    \\ \text{O} \end{array}$
W	$\begin{array}{c} \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{C} - \text{O} - \text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3 \\    \\ \text{O} \end{array}$

<i>Reagent 1</i>	$\text{SOCl}_2 / \text{PCl}_3 / \text{PCl}_5$
<i>Reagent 2</i>	conc. $\text{H}_2\text{SO}_4$