

# 3

91605



916050



NEW ZEALAND QUALIFICATIONS AUTHORITY  
MANA TOHU MĀTAURANGA O AOTEAROA

QUALIFY FOR THE FUTURE WORLD  
KIA NOHO TAKATŪ KI TŌ ĀMUA AO!

SUPERVISOR'S USE ONLY

## Level 3 Biology, 2015

### 91605 Demonstrate understanding of evolutionary processes leading to speciation

2.00 p.m. Monday 23 November 2015  
Credits: Four

Achievement	Achievement with Merit	Achievement with Excellence
Demonstrate understanding of evolutionary processes leading to speciation.	Demonstrate in-depth understanding of evolutionary processes leading to speciation.	Demonstrate comprehensive understanding of evolutionary processes leading to speciation.

Check that the National Student Number (NSN) on your admission slip is the same as the number at the top of this page.

**You should attempt ALL the questions in this booklet.**

If you need more room for any answer, use the extra space provided at the back of this booklet and clearly number the question.

Check that this booklet has pages 2–12 in the correct order and that none of these pages is blank.

**YOU MUST HAND THIS BOOKLET TO THE SUPERVISOR AT THE END OF THE EXAMINATION.**

**TOTAL**

ASSESSOR'S USE ONLY

## QUESTION ONE

‘Land lobsters’ are the common name of many species of large, flightless, ground-dwelling insects distributed in New Guinea, New Caledonia, and Lord Howe Island. Land lobsters have a stocky body form. Some males have enlarged and powerfully armed hind legs, and the females have an elongated ovipositor which they use to deposit eggs into the soil. Nuclear and mitochondrial DNA sequence analysis has shown that the different land lobsters species are unrelated to each other, and therefore have undergone convergent evolution.

*For copyright reasons,  
this resource cannot be  
reproduced here.*

Different ‘land lobster’ species, (a) to (f), compared with a winged, canopy-dwelling stick insect, (g).

Adapted from Buckley, T.E. et al. (2009). Extreme convergence in stick insect evolution: phylogenetic placement of the Lord Howe Island tree lobster. *Proc. R. Soc.* 276, 1055–1062.

Pōhutukawa (*Metrosideros excelsa*), northern rātā (*Metrosideros robusta*), and southern rātā (*Metrosideros umbellata*) are all related species belonging to the same genus. These species have undergone divergent evolution during the ice age that occurred between one and two million years ago.

Pōhutukawa has a coastal distribution and is very salt-tolerant. It has multiple trunks, is a coloniser of coastal cliffs and bare volcanic lava, and is susceptible to light frosts.

Northern rātā usually begins life as an epiphyte perched high on another tree. From here it sends down roots to form a trunk that can grow into a 40 m tree. It has moderate frost tolerance.

Southern rātā usually grows from the ground to a 15 m high, single-trunked tree that can tolerate frost and colder climates.

*For copyright reasons,  
this resource cannot be  
reproduced here.*

Different forms of *Metrosideros*.

Adapted from: P. Simpson, *Pohutukawa and Rata*, (Wellington, Te Papa Press, 2005), p. 125.

Discuss the evolutionary patterns AND selection pressures that have contributed to these patterns for land lobsters and *Metrosideros*.

In your answer:

- describe convergent evolution and divergent evolution
- explain, using the evidence given above, how each of these patterns could arise
- explain, by giving examples from the resource material, which pattern is associated with homologous structures AND which pattern is associated with analogous structures
- discuss why land lobsters have a different evolutionary pattern to *Metrosideros*.

ASSESSOR'S  
USE ONLY

**There is more space for your answer to this question on the following page.**




**This page has been deliberately left blank.**  
**The examination continues on the following page.**

## QUESTION TWO

The green anole lizard (*Anolis carolinensis*) is the only native anole in the United States. However, since 1940, the Cuban brown anole lizard (*Anolis sagrei*) has been invading the southeastern United States so that both species exist sympatrically in this area. Both species have adhesive scales on their toe pads called lamellae, and are very similar in habitat use, ecology, and dietary preferences. Biologists studying these anole compared the height at which the green anole perched in trees in the presence AND absence of the Cuban brown anole, and their results are shown in Figure 1. Biologists also measured toe pad area and lamella number in the green anole in the presence AND absence of the Cuban brown anole, and their results are shown in Figure 2a and Figure 2b.

**Figure 1: Perch height of green anole lizard in presence and absence of Cuban brown anole lizard**




*For copyright reasons, this resource cannot be reproduced here.*

Green anole and Cuban brown anole lizards.


<http://davewelling.photoshelter.com/image/10000HVzOE-fE2lmQ>

Adapted from: Stuart, Y. E., et al. (2014), 'Rapid evolution of a native species following invasion by a congener', *Science* 346 (6208): 463–466

**Figure 2a: Toe pad area in green anole lizard in absence and presence of Cuban brown anole lizard**



**Figure 2b: Lamella number in green anole lizard in absence and presence of Cuban brown anole lizard**



*For copyright reasons, this resource cannot be reproduced here.*

Green anole hind foot showing toe pads.

[www.utexas.edu/mews/2014/10/23/anole-lizards-evolution-florida/](http://www.utexas.edu/mews/2014/10/23/anole-lizards-evolution-florida/)

Adapted from: Stuart, Y. E., et al. (2014), 'Rapid evolution of a native species following invasion by a congener', *Science* 346 (6208): 463–466

Discuss the natural selection pressures that have affected evolution in the green anole.

In your answer:

- describe natural selection and the trends shown by the resource material
- explain the type of natural selection occurring in the green anole
- evaluate the impact of competition on the evolution of the green anole.

ASSESSOR'S  
USE ONLY

**There is more space for your  
answer to this question on the  
following page.**





The four-wing saltbush (*Atriplex canescens*) is a shrub that has undergone polyploidy. It has a haploid number of nine chromosomes ( $n = 9$ ). Biologists studied four-wing saltbushes with different numbers of chromosomes. Each type of saltbush lives in a slightly different habitat depending on how much water is available. Biologists measured the width of the water transport system (called the xylem) in each type of saltbush, and the results are shown in the table below. The xylem can be blocked by air bubbles in drought conditions.

Source: Hao, G et al. 'Polyploidy enhances the occupation of heterogeneous environments through hydraulic related trade-offs in *Atriplex canescens* (Chenopodiaceae)'. *New Phytologist* (2013) 197: 970–978.

Discuss the implications of polyploidy on the evolution of the four-wing saltbush.

- describe polyploidy and describe why the four-wing saltbush polyploids are fertile
- explain how polyploid formation could occur in the four-wing saltbush
- discuss what processes need to occur for the polyploids to become separate species
- discuss how the change in structure of the polyploids may lead to speciation.

**There is more space for your answer to this question on the following page.**





**Extra paper if required.**  
**Write the question number(s) if applicable.**

ASSESSOR'S  
USE ONLY

QUESTION  
NUMBER

91605